Claims:

5

10

1. A process for preparing aryllithium compounds of the formulae (V) and (VI) and reacting them with suitable electrophiles to give compounds of the formulae (VII) and (VIII), aryl halides of the formula (I) are reacted with lithium metal to generate a lithium compound (II), this is used for deprotonating the aromatic substrate (III) or (IV), and is finally converted by addition of the electrophilic component into the target compound of the formula (VII) or (VIII) (equation 1),

Step 1: Generation of the base

15 Step 2: Deprotonation of the substrate

Step 3: Reaction of the aryllithium compound with an electrophile

5

10

15

20

25

(Equation 1)

where Ar is phenyl, alkyl-substituted phenyl, fluorine- or chlorinesubstituted phenyl, naphthyl-, alkyl-substituted naphthyl or is biphenyl,

Hal = fluorine, chlorine, bromine or iodine,

the radicals X_{1-4} are, independently of one another, either carbon, X_iR_i (i=1-4) can symbolize nitrogen, or two radicals X_iR_i which are adjacent or connected via a formal double bond can together be O (furans), S (thiophenes), NH or NR_i (pyrroles),

Z is, in the case of benzoidal aromatics, a group which activates the ortho position, for example CF₃, OCF₃, Cl, F, Oalkyl, Oaryl, Salkyl, Saryl, CH₂OH, CH₂OR, CH(OR)₂, CONR₂, NHR, NR₂, or in the case of heterocycles has the same meaning as R₁₋₄,

the radicals R₁₋₄ are substituents selected from the group consisting of hydrogen, methyl, primary, secondary or tertiary, cyclic or acyclic alkyl radicals having from 2 to 12 carbon atoms, substituted cyclic or acyclic alkyl groups, alkoxy, dialkylamino, alkylamino, arylamino, diarylamino, phenyl, substituted phenyl, alkylthio, diarylphosphino, dialkylphosphino, dialkylaminocarbonyl or diarylaminocarbonyl, monoalkylaminocarbonyl or monoarylaminocarbonyl, CO₂alkyl,

5

 CO_2 , 1-hydroxyalkyl, 1-alkoxyalkyl, fluorine or chlorine, CN or heteroaryl, where two adjacent radicals R_{1-4} can together correspond to a fused-on aromatic or aliphatic ring,

- and "Electrophile" is any electrophilic component which can be reacted with aryllithium compounds.
- The process as claimed in claim 1, wherein the compounds of the formula (III) or (IV) are selected from the group consisting of benzenes, furans, thiophenes, pyridines, pyridazines, pyrimidines, pyrazines, N-substituted pyrroles, benzofurans, indoles and naphthalenes.
- The process as claimed in claim 1 or 2, characterized in that the 3. electrophile used is a compound selected from the following group: 15 oxirane, substituted oxirane, azomethine, aryl or alkyl cyanate, nitroenolate, immonium salts, haloaromatics, aryl triflates, other aryl sulfonates, carbon dioxide, carbon monoxide, aldehydes, ketones, α, β -unsaturated aldehydes or ketones, ketenes, alkali metal or alkaline earth metal salts of carboxylic acids, aliphatic nitriles, 20 aromatic nitriles, amides, esters and alkylating agents and boron electrophiles of the formula BW3, where the radicals W are identical or different and are each C1-C6-alkoxy, fluorine, chlorine, bromine, iodine, N(C1-C6-alkyl)2 or S(C1-C5-alkyl), and silicon electrophiles of the formula SiW4, where the radicals W are identical or different and 25 are each C₁-C₆-alkoxy, fluorine, chlorine, bromine, iodine, N(C₁-C₆ $alkyl)_2$ or $S(C_1-C_5-alkyl)$.
- 4. The process as claimed in at least one of the preceding claims, characterized in that the reaction is carried out in an organic ether solvent.
- 5. The process as claimed in at least one of the preceding claims, characterized in that the reaction temperature is in the range from -100 to +35°C.
 - 6. The process as claimed in at least one of the preceding claims, characterized in that the concentrations of the aliphatic or aromatic

intermediates of the formula (II) or (IV) are in the range from 5 to 30% by weight.

- 7. The process as claimed in at least one of the preceding claims, characterized in that the amount of lithium added per mole of halogen reacted is from 1.95 to 2.5 mol.
- 8. The process as claimed in at least one of the preceding claims, characterized in that organic redox systems are added to the reaction mixture, preferably in amounts of < 0.5 mol%.



onal Application No PCT/EP 03/09252

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 C07F1/02

According to International Patent Classification (IPC) or to both national classification and IPC

Minimum documentation searched (classification system followed by classification symbols) $IPC\ 7\ C07F$

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, BEILSTEIN Data

	INTS CONSIDERED TO BE RELEVANT	Relevant to claim No.
Category °	Citation of document, with indication, where appropriate, of the relevant passages	
Χ,Υ	SCHLOSSER ET AL: "Displacement of Halogens" April 2002 (2002-04), ORGANOMETALLICS IN SYNTHESIS. A MANUAL, PART I, CHAPTER 4.1, PAGES 86-112 AND 314-324 XP002261337 page 86 -page 112; table 18	1-8
Y	SCHLOSSER ET AL: "Organometallics in Synthesis, A Manual passage", ORGANOMETALLICS IN SYNTHESIS. A MANUAL, XX, XX, PAGE(S) 223-247,341-347 XP002228264 Seiten 229, 238, 241	1-8
	-/	

X Further documents are listed in the continuation of box C.	χ Patent family members are listed in annex.
Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date daimed Date of the actual completion of the international search	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "8" document member of the same patent family Date of mailing of the international search report
13 November 2003	09/12/2003
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL – 2280 HV Rijswijk Tel. (+31–70) 340–2040, Tx. 31 651 epo nl, Fax: (+31–70) 340–3016	Authorized officer Richter, H



Intersponal Application No PCT/EP 03/09252

		PCI/EP 03/09252
	ation) DOCUMENTS CONSIDERED TO BE RELEVANT	
Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 00 64905 A (GISSOT ARNAUD ; JOST SYLVIE (FR); WAGNER ALAIN (FR); DESMURS JEAN R) 2 November 2000 (2000-11-02) Seite 7, Zeilen 9-11 example 1; table 1	1-8
Ρ,Υ	WO 03 033503 A (CLARIANT GMBH ;MEUDT ANDREAS (DE)) 24 April 2003 (2003-04-24) the whole document	1-8
(SICE, JEAN: "Preparation and Reactions of 2-Methylthiophene" J. AM. CHEM: SOC., vol. 75, 1975, pages 3697-3700, XP002261336 page 3699, column 1, line 30 - line 42	1-8
l		1
[
- [
j		
İ		·
ļ	·	
		ŀ
		·
	,	



Interior bonal Application No PCT/EP 03/09252

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
WO 0064905	A	02-11-2000	FR AU WO	2793792 A1 4302600 A 0064905 A2	24-11-2000 10-11-2000 02-11-2000
WO 03033503	Α	24-04-2003	DE WO	10150615 A1 03033503 A2	30-04-2003 24-04-2003